

BioLiNE® Application Recommendations

Applicable with BioLiNE® Gold and Emerald

Crop	Application Stages
Corn	Starter: 300 mL/ac (10 oz) Foliar V3-V6: 500 mL/ac (17 oz) Foliar VT: 500 mL/ac (17 oz)
Soybean	Starter: 300 mL/ac (10 oz) Foliar V1-V3: 500 mL/ac (17 oz) Foliar R1-R5: 500 mL/ac (17 oz)
Cereals	Starter: 300 mL/ac (10 oz) Foliar stage 4-6: 400 mL/ac (14 oz) Foliar stage 9-10: 400 mL/ac (14 oz) Foliar stage 10-11: 400 mL/ac (14 oz)
Potato	In-furrow: 1L/ac (32 oz) Herbicide Tank Mix: 500 mL/ac (17 oz) Fungicide Tank Mix: 250 mL/ac (8 oz) every two weeks
Pasture Grasses	Start of regrowth: 300 mL/ac (10 oz) Animal rotation: 300 mL/ac (10 oz) Tank mix/impregnate onto fertilizer: 300 mL/ac (10 oz)
Beans, Peas	Pre-plant: 500 - 750 mL/ac (16-24 oz) with pre-emergence herbicide At planting: 300 mL/ac (10 oz) Post Emergence: 250 mL/ac (8 oz) at rapid vegetative growth and flowering stages

Crop	Application Stages
Citrus and Orchards	Apply 500 mL/ac (17 oz) every 30 days
Small Berries & Fruits	Foliar: 250 mL/ac (8 oz) every 10-14 days Fertigation: 250 mL/ac (8 oz) every 2 weeks
Cucurbits (cucumber, pumpkin, melon)	Transplant: 125 mL/400 L (4 oz/100 gal) transplant solution Apply 500 mL/ac (17 oz) every 14 days
Cruciferous (broccoli, cabbage, cauliflower)	At transplant: 100 mL/400L (4 oz/100 gal) transplant solution Apply 500 mL/ac (17 oz) at 2 leaf, 5-6 leaf, 6-8 leaf, cupping, and heading
Leafy Vegetables	Fertigation: 250 mL/ac (8 oz) every 2 weeks
Cannabis	Foliar: Apply 500 mL/ac (17 oz) every 2-3 weeks starting at 6-leaf stage Fertigation: Apply 750 mL/ac (25 oz) every 2-3 weeks Hydroponic: Add 2 mL/L (2.5 oz/10 gal) of reservoir water



The Certified Biostimulant program (U.S.-based) is used to certify products that meet industry standards for efficacy, safety, and composition.



The CE certification (EU-based) shows that biostimulant products meet the EU safety, health, and environmental protection requirements and prove the effectiveness of the label claims.

#FirstclassFulvic